

NPS55HH73101A



# CONDITIONS UNDER WHICH A PENALTY FUNCTION ALGORITHM IS WELL DEFINED

by

James K. Hartman

October 1973

Approved for public release; distribution unlimited.



NAVAL POSTGRADUATE SCHOOL  
Monterey, California

Rear Admiral M. B. Freeman, USN  
Superintendent

M. U. Clauser  
Provost

ABSTRACT:

Conditions are given which ensure that a recently developed exact penalty function algorithm for nonlinear programming problems is well defined.

Prepared by:



## I. THE PROBLEM

We consider the nonlinear programming problem with inequality constraints

$$\begin{array}{lll} \text{NLP} & \text{maximize} & f(x) \\ & \text{subject to} & g_i(x) \leq 0 \quad i = 1, \dots, m \\ & & x \in E^n \end{array} \quad (1)$$

where  $f$  and  $g_i$  are continuously differentiable and for simplicity we assume that a constraint qualification holds. In reference [1] we developed a penalty function algorithm which involves alternate cycles of 1) maximizing a penalty function and 2) approximating optimal Lagrange multipliers. In the interests of brevity the reader is referred to [1] for a complete description of the procedure and its convergence properties. For the purposes of this report it is enough to recall that at each maximization cycle we have fixed  $k > 0 \in E^1$ ,  $\lambda > 0 \in E^m$  and we want to find  $x \in E^n$  to maximize the penalty function

$$P(x, k, \lambda) = f(x) - (1/k) \sum_{i=1}^m \lambda_i [\exp(kg_i(x)) - 1] \quad (2)$$

In [2] Gould and Howe show that under certain weak conditions if  $x^*$  is a local maximum for NLP(1) with optimal Lagrange multipliers  $\lambda^*$ , then there exists a  $k$  sufficiently large that  $P(x, k, \lambda^*)$  has a local maximum at  $x^*$ . Thus, if  $\lambda^*$  is known and  $k$  is chosen large enough, the function  $P(x, k, \lambda^*)$  is an exact penalty function. The algorithm in [1] attempts to determine the optimal Lagrange multipliers iteratively while simultaneously searching for  $x^*$ . Thus, it involves maximizing

$P(x,k,\lambda)$  for fixed  $\lambda \neq \lambda^*$ , and it is important to determine conditions under which this maximization can be accomplished. If, for the particular choice of  $\lambda$  being used at some cycle, the  $P(x,k,\lambda)$  function is unbounded (as a function of  $x$ ) then the algorithm is not well defined.

## II. CONDITIONS FOR A WELL DEFINED ALGORITHM

In this section we give two sets of conditions and prove that under either set of conditions the penalty function has a finite global maximum for any  $k > 0 \in E^1$  and any  $\lambda > 0 \in E^m$ .

Theorem 1: If  $f(x) \rightarrow -\infty$  as  $\|x\| \rightarrow \infty$  then the algorithm is well defined.

Proof: The proof is simple and requires only that we examine the penalty term for the  $i$ th constraint

$$-(1/k) \lambda_i [\exp(kg_i(x)) - 1] \quad (3)$$

with  $k > 0$ ,  $\lambda_i > 0$ . As  $g_i(x) \rightarrow \infty$  (infeasible), (3)  $\rightarrow -\infty$ ; but as  $g_i(x) \rightarrow -\infty$  (feasible) (3)  $\rightarrow \lambda_i/k$ .

Thus,

$$-(1/k) \sum_{i=1}^m \lambda_i \exp[(kg_i(x)) - 1] \leq \sum_{i=1}^m \lambda_i/k \quad (4)$$

So if  $f(x) \rightarrow -\infty$  for all  $x$  with  $\|x\| \rightarrow \infty$ , then also

$P(x, k, \lambda) \rightarrow -\infty$  as  $\|x\| \rightarrow \infty$ . Hence,  $P$  is not unbounded for maximization, and this is true for any  $k > 0 \in E^1$  and any  $\lambda \geq 0 \in E^m$ .  $\square$

Theorem 2: If a)  $f(x)$  is a strictly concave function

b)  $g_i(x)$  are convex functions for all  $i = 1, \dots, m$ , and

c) NLP(1) has finite optimal solution  $x^*$  with optimal Lagrange multipliers  $\lambda^*$

then for any fixed  $k > 0 \in E^1$ ,  $\lambda > 0 \in E^m$ ,  $\exists$  a finite  $x$  which maximizes  $P(x, k, \lambda)$ .

Proof: For fixed  $k > 0$ ,  $\lambda > 0$  we know that  $P(x, k, \lambda)$  is a strictly concave function of  $x$  because of assumptions a) and b). To show that a finite maximum exists it suffices to show that a finite maximum along any half line exists, that is,  $\forall x, d \in E^n$ , as  $\theta \in E^1 \rightarrow \infty$  the derivative

$$P'(\theta) = \frac{d}{d\theta} P(x + \theta d, k, \lambda) \quad (5)$$

must eventually become negative (or equal zero).

Since  $x^*$  maximizes  $P(x, k, \lambda^*)$  (see [2]) we know this is true for  $\lambda = \lambda^*$ .

Thus, consider any  $x, d \in E^n$ ,  $k > 0 \in E^1$ ,  $\lambda > 0 \in E^m$ .

$$\begin{aligned} P'(\theta) &= \frac{d}{d\theta} P(x + \theta d, k, \lambda) \\ &= \frac{d}{d\theta} \left\{ f(x + \theta d) - (1/k) \sum_{i=1}^m \lambda_i [\exp(k g_i(x + \theta d)) - 1] \right\} \quad (6) \\ &= f'(\theta) - \sum_{i=1}^m \left( \frac{\lambda_i}{k} \right) \exp[k g_i(\theta)] g_i'(\theta) \end{aligned}$$

where we have introduced the abbreviations  $f'(\theta) \equiv \frac{d}{d\theta} f(x + \theta d)$ ,

$g_i(\theta) \equiv g_i(x + \theta d)$ ,  $g_i'(\theta) \equiv \frac{d}{d\theta} g_i(x + \theta d)$ .



There are several cases for each of  $f$  and  $g_i$  :

Cases for  $f$  (concave)

f1  $f(\theta) \rightarrow -\infty$  as  $\theta \rightarrow \infty$

f2  $f(\theta)$  is non decreasing as  $\theta \rightarrow \infty$

then  $f'(\theta) > 0$  and decreasing as  $\theta \rightarrow \infty$  (7)

Cases for each  $g_i$  (convex)

g1  $g_i(\theta) \rightarrow \infty$  as  $\theta \rightarrow \infty$

then  $\exists r > 0$  such that eventually  $g_i'(\theta) \geq r$  so that

$$-\frac{\lambda_1}{k} [\exp kg_i(\theta)] g_i'(\theta) \rightarrow -\infty \text{ as } \theta \rightarrow \infty \quad (8)$$

g2  $g_i(\theta)$  is non increasing as  $\theta \rightarrow \infty$  then  $\exp kg_i(\theta) > 0$  and non increasing as  $\theta \rightarrow \infty$ ,  $g_i'(\theta) \leq 0$  and non decreasing as  $\theta \rightarrow \infty$ , so  $-\frac{\lambda_1}{k} [\exp kg_i(\theta)] g_i'(\theta) \geq 0$  and non increasing as  $\theta \rightarrow \infty$ . (9)

Analysis of Cases

1. If f1 holds, then  $P(\theta) \rightarrow -\infty$  as  $\theta \rightarrow \infty$  for any combination of cases on the  $g_i$ . (see Theorem 1) Thus,  $P$  has a finite global maximum for any  $\lambda > 0$ .
2. If f2 holds,
  - 2a) If also  $\exists i$  with  $g_i$  in case g1 then (7), (8), (9)

show that  $P'(\theta)$  in (6) approaches  $-\infty$  as  $\theta \rightarrow \infty$ .

Thus  $P'(\theta)$  must eventually become negative so that

$P$  has a finite max along the half line.

2b) Otherwise for all  $i = 1, \dots, m$  constraint  $g_i$  is in case g2.

then (6), (7) and (9) show that  $P'(\theta) > 0$  as  $\theta \rightarrow \infty$

so that no maximum is attained. But this is true for

any  $\lambda$  including  $\lambda = \lambda^*$ .

Thus,  $P(x, k, \lambda^*)$  does not have a finite maximum contrary

to assumption. Hence case 2b cannot occur if NLP

has a finite optimal solution. □.

### III. SUMMARY

Two sets of conditions which guarantee that the exact penalty function algorithm in [1] is well defined have been presented and proved. It should perhaps also be emphasized that the algorithm in [1] is primarily a method for locating local solutions to NLP (1), and that it can frequently do this even if  $P(x,k,\lambda)$  is globally unbounded, (for non concave problems) since in the region of search  $P(x,k,\lambda)$  will still have a local maximum. In such a case, however, the possibility of straying into the global unbounded region cannot be overlooked.

## REFERENCES

- [1] J. K. Hartman, "Iterative Determination of Parameters for an exact Penalty Function". Naval Postgraduate School Technical Report NPS55HH71121A, Dec. 1971.
- [2] F. J. Gould and S. Howe, "A New Result on Interpreting Lagrange Multipliers as Dual Variables". University of North Carolina Institute of Statistics Mimeo Series No. 738, January 1971.

## INITIAL DISTRIBUTION LIST

Defense Documentation Center Cameron Station Alexandria, Virginia 22314	12
Dean of Research Code 023 Naval Postgraduate School Monterey, California 93940	1
Library, Code 0212 Naval Postgraduate School Monterey, California 93940	2
Dr. Thomas Varley Office of Naval Research Arlington, Virginia 22217	1
Library, Code 55 Department of Operations Research and Administrative Sciences Naval Postgraduate School Monterey, California 93940	5
Professor Donald P. Gaver Department of Operations Research and Administrative Sciences Naval Postgraduate School Monterey, California 93940	1
Professor G. T. Howard Department of Operations Research and Administrative Sciences Naval Postgraduate School Monterey, California 93940	1
Professor James K. Hartman Department of Operations Research and Administrative Sciences Naval Postgraduate School Monterey, California 93940	10

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS55HH73101A	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  CONDITIONS UNDER WHICH A PENALTY FUNCTION ALGORITHM IS WELL DEFINED		5. TYPE OF REPORT & PERIOD COVERED  Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  James K. Hartman		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS  Naval Postgraduate School Monterey, California 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE October 1973
		13. NUMBER OF PAGES 13
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Nonlinear Programming, Penalty Functions, Mathematical Programming Optimization, Lagrange Multipliers		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  Conditions are given which ensure that a recently developed exact penalty function algorithm for nonlinear programming problems is well defined.		

U158248

DUDLEY KNOX LIBRARY - RESEARCH REPORTS



5 6853 01058001 2

~~U1582~~